

Levy Processes, Generators

Sakhnovich Lev

E-mail address: lev.sakhnovich@verizon.net

Abstract. For a broad class of the Levy processes the new form (convolution type) of the infinitesimal generators is introduced. It leads to the new notions: a truncated generator, a quasi-potential. The probability of the Levy process remaining within the given domain is estimated.

1 Main notions

Let us consider the Levy processes X_t on R . If $P(X_0 = 0) = 1$ then Levy-Khinchine formula gives [1],[6]

$$\mu(x, t) = E\{\exp[izX_t]\} = \exp[-t\lambda(z)], \quad t \geq 0, \quad (1.1)$$

where

$$\lambda(z) = \frac{1}{2}Az^2 - i\gamma z - \int_{-\infty}^{\infty} (e^{izx} - 1 - izx1_{D(x)})d\nu(x). \quad (1.2)$$

Here $A \geq 0$, $\gamma = \overline{\gamma}$, and $D = \{x : |x| \leq 1\}$ is closed unit ball, $\nu(x)$ is monotonically increasing function satisfying the conditions

$$\int_{-\infty}^{\infty} \frac{x^2}{1+x^2} d\nu(x) < \infty. \quad (1.3)$$

By $P_t(x_0, B)$ we denote the probability $P(X_t \in B)$ when $P(X_0 = x_0) = 1$ and $B \in R$. The transition operator is defined by the formula

$$P_t f(x) = \int_{-\infty}^{\infty} P_t(x, dy) f(y). \quad (1.4)$$

Let C_0 be the Banach space of continuous functions $f(x)$ ($-\infty < x < \infty$) satisfying the condition $\lim_{|x| \rightarrow \infty} f(x) = 0$, with the norm $\|f\| = \sup_x |f(x)|$. We denote by C_0^n the set of $f(x) \in C_0$ such that $f^{(k)}(x) \in C_0$, ($1 \leq k \leq n$). It is known that [6]

$$P_t f \in C_0 \quad (1.5)$$

if $f(x) \in C_0$.

Now we formulate the following important result (see [2],[6]) .

Theorem 1.1. *The family of the operators P_t ($t \geq 0$) defined by the Levy process X_t is a strong continuous semigroup on C_0 with norm $\|P_t\| = 1$. Let L be its infinitesimal generator. Then*

$$Lf = \frac{1}{2}A \frac{d^2 f}{dx^2} + \gamma \frac{df}{dx} + \int_{-\infty}^{\infty} (f(x+y) - f(x) - y \frac{df}{dx} 1_{D(x)}) d\nu(x), \quad (1.6)$$

where $f \in C_0^2$.

2 Convolution type form of infinitesimal generator

1. In this section we prove that under some conditions the infinitesimal generator L can be represented in the special convolution type form

$$Lf = \frac{d}{dx} S \frac{d}{dx} f, \quad (2.1)$$

where the operator S is defined by the relation

$$Sf = \frac{1}{2}Af + \int_{-\infty}^{\infty} k(y-x)f(y)dy, \quad (2.2)$$

and for arbitrary M ($0 < M < \infty$) we have

$$\int_{-M}^M |k(t)|dt < \infty. \quad (2.3)$$

The representation L in form (2.1) is convenient as the operator L is expressed with the help of the classic differential and convolution operators.

By C_c we denote the set of functions $f(x) \in C_0$ with compact support.

Lemma 2.1. *Let the following conditions be fulfilled.*

1. *The function $\nu(x)$ is monotonically increasing, has the derivative when $x \neq 0$ and*

$$\int_{-\infty}^{\infty} \frac{x^2}{1+x^2} d\nu(x) = \int_{-\infty}^{\infty} \frac{x^2}{1+x^2} \nu'(x) dx < \infty, \quad (2.4)$$

$$\nu(x) \rightarrow 0, \quad x \rightarrow \infty. \quad (2.5)$$

2. For arbitrary $M(0 < M < \infty)$ we have

$$\int_{-M}^M |\nu(x)| dx < \infty, \quad \int_{-M}^M |x| \nu'(x) dx < \infty. \quad (2.6)$$

3.

$$x\nu(x) \rightarrow 0, \quad x \rightarrow 0. \quad (2.7)$$

Then the equality

$$J = \int_{-\infty}^{\infty} [f(y+x) - f(x)] \nu'(y) dy, \quad (2.8)$$

is true, where $f(x) \in C_0^2$, $k(x) = \int_0^x \nu(y) dy$ and

$$J = \frac{d}{dx} \int_{-\infty}^{\infty} f'(y) k(y-x) dy. \quad (2.9)$$

Proof. For every $f(x) \in C_c$ there exists such $M(0 < M < \infty)$ that

$$f(x) = 0, \quad x \notin [-M, M]. \quad (2.10)$$

Let us introduce the following notations

$$J_1 = \frac{d}{dx} \int_{-\infty}^x f'(y) k(y-x) dy, \quad (2.11)$$

$$J_2 = \frac{d}{dx} \int_x^{\infty} f'(y) k(y-x) dy. \quad (2.12)$$

Using (2.11) we have

$$J_1 = -\frac{d}{dx} \int_{-M}^x [f(y) - f(x) + f(x)] k'(y-x) dy. \quad (2.13)$$

From (2.11) and (2.13) we deduce the relation

$$J_1 = f(x) k'(-M-x) + \int_{-M}^x [f(y) - f(x)] k''(y-x) dy. \quad (2.14)$$

When $M \rightarrow \infty$ we obtain the equality

$$J_1 = \int_{-\infty}^0 [f(y+x) - f(x)] k''(y) dy. \quad (2.15)$$

In the same way we deduce the relation

$$J_2 = \int_0^\infty [f(y+x) - f(x)]k''(y)dy. \quad (2.16)$$

The relation (2.9) follows directly from (2.15) , (2.16) and equality $J = J_1 + J_2$. The lemma is proved.

Lemma 2.2. *Let the following conditions be fulfilled.*

1. *The function $\nu(x)$ satisfies conditions of Lemma 2.1.*
2. *For arbitrary M ($0 < M < \infty$) we have*

$$\int_{-M}^M |k(x)|dx < \infty, \quad \int_{-M}^M |x\nu(x)|dx < \infty, \quad (2.17)$$

where

$$k'(x) = \nu(x), \quad x \neq 0. \quad (2.18)$$

3.

$$xk(x) \rightarrow 0, \quad x \rightarrow 0; \quad x^2\nu(x) \rightarrow 0, \quad x \rightarrow 0. \quad (2.19)$$

Then the equality

$$J = \int_{-\infty}^\infty [f(y+x) - f(x) - y \frac{df(x)}{dx} 1_{D(y)}] \nu'(y) dy + \Gamma f'(x), \quad (2.20)$$

is true, where $\Gamma = \bar{\Gamma}$ and $f(x) \in C_c$.

Proof. From (2.11) we obtain the relation

$$J_1 = f'(x)\gamma_1 - \int_{x-1}^x [f'(y) - f'(x)]k'(y-x)dy - \int_{-M}^{x-1} f'(y)k'(y-x)dy, \quad (2.21)$$

where $\gamma_1 = k(-1)$. We introduce the notations

$$P_1(x, y) = f(y) - f(x) - (y-x)f'(x), \quad P_2(x, y) = f(y) - f(x). \quad (2.22)$$

Integrating by parts (2.21) and passing to limit when $M \rightarrow \infty$ we deduce that

$$J_1 = f'(x)\gamma_2 + \int_{x-1}^x P_1(x, y)k''(y-x)dy + \int_{-M}^{x-1} P_2(x, y)k''(y-x)dy, \quad (2.23)$$

where $\gamma_2 = k(-1) - k'(-1)$. It follows from (2.22) and (2.23) that

$$J_1 = \int_{-\infty}^x [f(y+x) - f(x) - y \frac{df(x)}{dx} 1_{D(y)}] \nu'(y) dy + \gamma_2 f'(x). \quad (2.24)$$

In the same way can be proved the relation

$$J_2 = \int_x^{\infty} [f(y+x) - f(x) - y \frac{df(x)}{dx} 1_{D(y)}] \nu'(y) dy + \gamma_3 f'(x), \quad (2.25)$$

where $\gamma_3 = -k(1) + k'(1)$. The relation (2.20) follows directly from (2.24) and (2.25). Here $\Gamma = \gamma_2 + \gamma_3$. The lemma is proved.

Remark 2.1. The operator $L_0 f = \frac{df}{dx}$ can be represented in form (2.1), where

$$S_0 f = \int_{-\infty}^{\infty} p_0(y-x) f(y) dy, \quad (2.26)$$

$$p_0(x) = \frac{1}{2} \text{sign}(x). \quad (2.27)$$

From Lemmas 2.1 , 2.2 and Remark 2.1 we deduce the following assertion.

Theorem 2.1. *Let the conditions of Lemma 2.1 or Lemma 2.2 be fulfilled . Then the corresponding operator L has form (2.1),(2.2).*

Proposition 2.1. *The generator L of the Levy process X_t admits the convolution representation (2.1), (2.2) if there exist such $C > 0$ and $0 < \alpha < 2$, $\alpha \neq 1$ that*

$$\nu'(y) \leq C |y|^{-\alpha-1}, \quad (2.28)$$

Proof. The function $\nu(y)$ has the form

$$\nu(y) = \int_{-\infty}^y \nu'(t) dt 1_{y < 0} - \int_y^{\infty} \nu'(t) dt 1_{y > 0}. \quad (2.29)$$

We suppose that $1 < \alpha < 2$ and introduce the function

$$k_0(y) = \int_{-\infty}^y (y-t) \nu'(t) dt 1_{y < 0} - \int_y^{\infty} (t-y) \nu'(t) dt 1_{y > 0}. \quad (2.30)$$

We obtain the relation

$$k(y) = k_0(y) + (\gamma - \Gamma) p_0(y), \quad (2.31)$$

where $k_0(y)$ and $p_0(y)$ are defined by (2.27) and (2.30) respectively. The constant Γ is defined by relations:

$$\Gamma = k_0(-1) - k'_0(-1) - k_0(-1) + k'_0(1), \quad 1 \leq \alpha < 2, \quad (2.32)$$

It follows from (2.28)-(2.30) that the conditions of Theorem 2.1 are fulfilled. Hence the proposition is true when $1 < \alpha < 2$. Let us consider the case when $0 < \alpha < 1$. In this case we have

$$k_0(y) = \int_{-\infty}^y \nu'(t) dt y + \int_y^0 \nu'(t) dt], \quad y < 0, \quad (2.33)$$

$$k_0(y) = - \int_y^{\infty} \nu'(t) dt y - \int_0^y \nu'(t) dt], \quad y > 0, \quad (2.34)$$

and

$$k(y) = k_0(y) + \gamma p_0(y) \quad 0 < \alpha < 1. \quad (2.35)$$

In view of (2.28) and (2.33),(2.34) the conditions of Theorem 2.1 are fulfilled. Hence the proposition is proved.

Corollary 2.1. *If condition (2.28) is fulfilled then*

$$k_0(y) \geq 0, \quad -\infty < y < \infty, \quad 1 < \alpha < 2, \quad (2.36)$$

$$k_0(y) \leq 0, \quad -\infty < y < \infty, \quad 0 < \alpha < 1. \quad (2.37)$$

Example 2.1. The stable processes.

For the stable processes we have $A = 0$, $\gamma = \bar{\gamma}$ and

$$\nu'(y) = |y|^{-\alpha-1} (C_1 1_{y<0} + C_2 1_{y>0}), \quad (2.38)$$

where $C_1 > 0$ $C_2 > 0$. Hence the function $\nu(y)$ has the form

$$\nu(y) = \frac{1}{\alpha} |y|^{-\alpha} (C_1 1_{y<0} - C_2 1_{y>0}). \quad (2.39)$$

Let us introduce the functions

$$k_0(y) = \frac{1}{\alpha(\alpha-1)} |y|^{1-\alpha} (C_1 1_{y<0} + C_2 1_{y>0}), \quad (2.40)$$

where $0 < \alpha < 2$, $\alpha \neq 1$. When $\alpha = 1$ we have

$$k_0(y) = -\log|y| (C_1 1_{y<0} + C_2 1_{y>0}). \quad (2.41)$$

It means that the conditions of proposition 2.1 are fulfilled. Hence the generator L for the stable processes admits the convolution type representation (2.1),(2.2).

Proposition 2.2. *The kernel $k(y)$ of the operator S in representation (2.1) for the stable processes has form (2.31), when $1 \leq \alpha < 2$, and has form (2.35) when $0 < \alpha < 1$.*

Example 2.2. The variance damped Levy processes [7].

For the variance damped Levy processes we have $A = 0$, $\gamma = \bar{\gamma}$ and

$$\nu'(y) = C_1 e^{-\lambda_1 |y|} |y|^{-\alpha-1} 1_{y < 0} + C_2 e^{-\lambda_2 |y|} |y|^{-\alpha-1} 1_{y > 0}, \quad (2.42)$$

where $C_1 > 0$, $C_2 > 0$, $\lambda_1 > 0$, $\lambda_2 > 0$, $0 < \alpha < 2$, $\alpha \neq 1$. It follows from (2.42) that the conditions of Proposition 2.1 are fulfilled. Hence the generator L for the variance damped Levy processes admits the convolution type representation (2.1),(2.2) and the kernel $k(y)$ is defined by formulas (2.30),(2.31), when $1 \leq \alpha < 2$, and by formula (2.35) when $0 < \alpha < 1$.

Example 2.3. The variance Gamma process [7].

For the variance Gamma process we have $A = 0$, $\gamma = \bar{\gamma}$ and

$$\nu'(y) = C_1 e^{-G|y|} |y|^{-1} 1_{y < 0} + C_2 e^{-M|y|} |y|^{-1} 1_{y > 0}, \quad (2.43)$$

where $C_1 > 0$, $C_2 > 0$, $G > 0$, $M > 0$. It follows from (2.43) that the conditions of Proposition 2.1 are fulfilled and the generator L of variance Gamma process admits the convolution type representation (2.1),(2.2). The kernel $k(y)$ is defined by relations (2.33) and (2.34).

Example 2.4. The normal inverse Gaussian process [7].

In the case of the normal inverse Gaussian process we have $A = 0$, $\gamma = \bar{\gamma}$ and

$$\nu'(y) = C e^{\beta y} K_1(|y|) |y|^{-1}, \quad C > 0, \quad -1 \leq \beta \leq 1, \quad (2.44)$$

where $K_\lambda(x)$ denotes the modified Bessel function of the third kind with index λ . Using equalities

$$|K_1(|x|)| \leq M e^{-|x|} / |x|, \quad M > 0, \quad 0 < x_0 \leq |x|, \quad (2.45)$$

$$|K_1(|x|)x| \leq M, \quad 0 \leq |x| \leq x_0 \quad (2.46)$$

we see that the conditions of Proposition 2.1 are fulfilled. Hence the corresponding generator L admits the convolution type representation (2.1, 2.2)

and the kernel $k(y)$ is defined by relations (2.33) and (2.34).

Example 2.5. The Meixner process [7].

For the Meixner process we have

$$\nu'(y) = C \frac{\exp \beta x}{x \sinh \pi x}, \quad (2.47)$$

where $C > 0$, $-\pi < \beta < \pi$. The conditions of Proposition 2.1 are fulfilled. Hence the corresponding generator L admits the convolution type representation (2.1),(2.2) and the kernel $k(y)$ is defined by relations (2.33),(2.34).

Remark 2.1. Examples 2.1-2.5 are used in the finance problems [7].

Example 2.6.Compound Poisson process.

We consider the case when $A = 0$, $\gamma = 0$ and

$$M = \int_{-\infty}^{\infty} \nu'(y) dy < \infty. \quad (2.48)$$

Using formulas (2.1) and (2.2) we deduce that the corresponding generator L has the following form

$$Lf = -Mf(x) + \int_{-\infty}^{\infty} \nu'(y-x)f(y)dy. \quad (2.49)$$

3 Potential

The operator

$$Qf = \int_0^{\infty} (P_t f) dt \quad (3.1)$$

is called *potential* of the semigroup P_t . The generator L and the potential Q are (in general) unbounded operators. Therefore the operators L and Q are defined not in the whole space $L^2(-\infty, \infty)$ but only in the subsets D_L and D_Q respectively. We use the following property of potential Q (see[6]).

Proposition 3.1. If $f = Qg$, $(g \in D_Q)$ then $f \in D_L$ and

$$-Lf = g. \quad (3.2)$$

Example 3.1. Compound Poisson process.

Let the generator L has form (2.49) where

$$M = \int_{-\infty}^{\infty} \nu'(x) dx < \infty, \quad \int_{-\infty}^{\infty} [\nu'(x)]^2 dx < \infty. \quad (3.3)$$

We introduce the functions

$$K(u) = -\frac{1}{M\sqrt{2\pi}} \int_{-\infty}^{\infty} \nu'(x) e^{-iux} dx, \quad (3.4)$$

$$N(u) = \frac{K(u)}{1 - \sqrt{2\pi}K(u)}. \quad (3.5)$$

Let us note that

$$|K(u)| < \frac{1}{\sqrt{2\pi}}, \quad u \neq 0; \quad K(0) = -\frac{1}{\sqrt{2\pi}}. \quad (3.6)$$

It means that $N(u) \in L^2(-\infty, \infty)$. Hence the function

$$n(x) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} N(u) e^{-iux} du \quad (3.7)$$

belongs to $L^2(-\infty, \infty)$ as well. It follows from (2.49), (3.2) and (3.7) that the corresponding potential Q has form (see [8], Ch.11)

$$Qf = \frac{1}{M} [f(x) + \int_{-\infty}^{\infty} f(y) n(x-y) dy]. \quad (3.8)$$

Proposition 3.2. *Let conditions (3.3) be fulfilled. Then the operators L and Q are bounded in the space $L^2(-\infty, \infty)$.*

Now we shall give an example when the kernel $n(x)$ can be written in an explicit form.

Example 3.2. We consider the case when

$$\nu'(x) = e^{-|x|}, \quad -\infty < x < \infty. \quad (3.9)$$

In this case $M = 2$ and the operator L takes the form

$$Lf = -2f(x) + \int_{-\infty}^{\infty} f(y) e^{-|x-y|} dy. \quad (3.10)$$

Formulas (3.4)-(3.7) imply that

$$Qf = \frac{1}{2}f(x) - \frac{1}{4\sqrt{2}} \int_{-\infty}^{\infty} f(y) e^{-|x-y|\sqrt{2}} dy. \quad (3.11)$$

4 Truncated generators and quasi-potentials

Let us denote by Δ the set of segments $[a_k, b_k]$ such that $a_1 < b_1 < a_2 < b_2 < \dots < a_n < b_n$, $1 \leq k \leq n$. By C_Δ we denote the set of functions $g(x)$ on $L^2(\Delta)$ such that

$$g(a_k) = g(b_k) = g'(a_k) = g'(b_k), \quad 1 \leq k \leq n, \quad g''(x) \in L^2(\Delta). \quad (4.1)$$

We introduce the operator P_Δ by relation $P_\Delta f(x) = f(x)$ if $x \in \Delta$ and $P_\Delta f(x) = 0$ if $x \notin \Delta$.

Definition 4.1. The operator

$$L_\Delta = P_\Delta L P_\Delta \quad (4.2)$$

is called a *truncated generator*.

Definition 4.2. The operator B with dense in $L^2(\Delta)$ definition domain L_B is called a *quasi-potential* if the functions $f = Bg$, ($g \in L_B$) belong to C_Δ and

$$-L_\Delta f = g. \quad (4.3)$$

It follows from (4.3) that

$$-P_\Delta L f = g, \quad (f = Bg). \quad (4.4)$$

Remark 4.1. In a number of cases (see the next section) we need relation (4.4). In these cases we can use the quasi-potential B , which is often simpler than the corresponding potential Q .

Remark 4.2. The operators of type (4.2) are investigated in book ([5], Ch.2).

Definition 4.3. We call the operator B a *regular* one if the following conditions are fulfilled.

1). The operator B has the form

$$Bf = \int_\Delta \Phi(x, y) f(y) dy, \quad f(y) \in L^2(\Delta), \quad (4.5)$$

where the function $\Phi(x, y)$ can have a discontinuity only when $x = y$ and

$$|\Phi(x, y)| \leq M |x - y|^{-\beta}, \quad 0 \leq \beta < 1. \quad (4.6)$$

2). The range of B is dense in $L^2(\Delta)$.

It follows from condition 2) and relation (4.4) that the range of L is dense in $L^2(\Delta)$. Further we assume that the quasi-potential B is regular.

5 The Probability of the Levy process remaining within the given domain

In many theoretical and applied problems it is important to estimate the quantity

$$p(t, \Delta) = P\{X_\tau \in \Delta\}, \quad 0 \leq \tau \leq t, \quad (5.1)$$

i.e. the probability that a sample of the process X_τ remains inside Δ for $0 \leq \tau \leq t$.

To derive the integro-differential equations corresponding to Levy processes we use the argumentation by Kac [3] and our own argumentation (see [5]). Now we get rid of the requirement for the process to be stable.

We suppose that

$$M(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} |\mu(x, t)| dx < \infty, \quad t > 0 \quad (5.2)$$

and

$$\int_0^1 M(t) dt < \infty. \quad (5.3)$$

According to (1.2) we have the inequality

$$\operatorname{Re}[\lambda(x)] \geq 0. \quad (5.4)$$

Remark 5.1. It follows from (5.2) and (5.4) that

$$\int_{-\infty}^{\infty} |\mu(x, t)|^p dx < \infty, \quad t > 0, \quad p \geq 1. \quad (5.5)$$

Proposition 5.1. (see [], Ch.5.) *Let condition (5.2) be fulfilled. Then the corresponding Levy process has the continuous density*

$$\rho(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-ixz} \mu(z, t) dz, \quad t > 0 \quad (5.6)$$

and

$$\rho(x, t) \leq M(t). \quad (5.7)$$

Now we introduce the sequence of functions

$$Q_{n+1}(x, t) = \int_0^t \int_{-\infty}^{\infty} Q_0(x - \xi, t - \tau) V(\xi) Q_n(\xi, \tau) d\xi d\tau, \quad (5.8)$$

where

$$Q_0(x, t) = \rho(x, t), \quad V(x) = 1 - 1_\Delta. \quad (5.9)$$

For Levy processes the following relation

$$Q_0(x, t) = \int_{-\infty}^{\infty} Q_0(x - \xi, t - \tau) Q_0(\xi, \tau) d\xi \quad (5.10)$$

is true. Using (5.8) and (5.10) we have

$$0 \leq Q_n(x, t) \leq t^n Q_0(x, t) / n!. \quad (5.11)$$

Hence the series

$$Q(x, t, u) = \sum_{n=0}^{\infty} (-1)^n u^n Q_n(x, t) \quad (5.12)$$

converges. The probabilistic meaning of $Q(x, t, u)$ is defined by the relation (see [4], Ch.4)

$$E\{\exp[-u \int_0^t V(X(\tau)) d\tau], c_1 < X(t) < c_2\} = \int_{c_1}^{c_2} Q(x, t, u) dx. \quad (5.13)$$

The inequality $V(x) \geq 0$ and relation (5.13) imply that the function $Q(x, t, u)$ monotonically decreases with respect to the variable "u" and the formulas

$$0 \leq Q(x, t, u) \leq Q(x, t, 0) = Q_0(x, t) = \rho(x, t) \quad (5.14)$$

are true. In view of (5.6), (5.7) and (5.14) the Laplace transform

$$\psi(x, s, u) = \int_0^{\infty} e^{-st} Q(x, t, u) dt, \quad s > 0. \quad (5.15)$$

has the meaning. According to (5.8) the function $Q(x, t, u)$ is the solution of the equation

$$Q(x, t, u) + u \int_0^t \int_{-\infty}^{\infty} \rho(x - \xi, t - \tau) V(\xi) Q(\xi, \tau, u) d\xi d\tau = \rho(x, t) \quad (5.16)$$

Taking from both parts of (5.16) the Laplace transform and bearing in mind (5.15) we obtain

$$\psi(x, s, u) + u \int_{-\infty}^{\infty} V(\xi) R(x - \xi, s) \psi(\xi, s, u) d\xi = R(x, s), \quad (5.17)$$

where

$$R(x, s) = \int_0^\infty e^{-st} \rho(x, t) dt. \quad (5.18)$$

Multiplying both parts (5.17) by $\exp(ixp)$ and integrating them with respect to x ($-\infty < x < \infty$) we have

$$\int_{-\infty}^\infty \psi(x, s, u) e^{ixp} [s + \lambda(p) + uV(x)] dx = 1. \quad (5.19)$$

Here we use relations (1.1), (5.6) and (5.18). Now we introduce the function

$$h(p) = \frac{1}{2\pi} \int_\Delta e^{-ixp} f(x) dx, \quad (5.20)$$

where the function $f(x)$ belongs to C_Δ . Multiplying both parts of (5.19) by $h(p)$ and integrating them with respect to p ($-\infty < p < \infty$) we deduce the equality

$$\int_{-\infty}^\infty \int_{-\infty}^\infty \psi(x, s, u) e^{ixp} [s + \lambda(p)] h(p) dx dp = f(0). \quad (5.21)$$

We have used the relations

$$V(x)f(x) = 0, \quad -\infty < x < \infty, \quad (5.22)$$

$$\frac{1}{2\pi} \lim_{N \rightarrow \infty} \int_{-N}^N \int_\Delta e^{-ixp} f(x) dx dp = f(0), \quad (5.23)$$

Since the function $Q(x, t, u)$ monotonically decreases with respect to " u " then by (5.15) this is true for the function $\psi(x, s, u)$ also. Hence there exists the limit

$$\psi(x, s) = \lim_{u \rightarrow \infty} \psi(x, s, u), \quad (5.24)$$

where

$$\psi(x, s) = 0, \quad x \notin \Delta. \quad (5.25)$$

The probabilistic meaning of $\psi(x, s)$ follows from the equality

$$\int_{-\infty}^\infty e^{-st} p(t, \Delta) dt = \int_\Delta \psi(x, s) dx. \quad (5.26)$$

Using the properties of the Fourier transform and conditions (5.24) , (5.25) we deduce from (5.21) the following assertion.

Proposition 5.2. *Let conditions (5.2) and (5.3) be fulfilled. Then the relation*

$$((sI - L_\Delta)f, \psi(x, s))_\Delta = f(0) \quad (5.27)$$

is true.

The important function $\psi(x, s)$ can be expressed with the help of the quasi-potential B .

Theorem 5.1. *Let conditions (5.2), (5.3) be fulfilled and let the quasi-potential be regular. Then in the space $L(\Delta)$ there is one and only one function*

$$\psi(x, s) = (I + sB^*)^{-1}\Phi(0, x) \quad (5.28)$$

which satisfies relation (5.27).

Proof. In view of (4.4) we have

$$-BL_\Delta f = f, \quad f \in C_\Delta. \quad (5.29)$$

Relations (5.28) and (5.29) imply that

$$((sI - L_\Delta)f, \psi(x, s))_\Delta = -((I + sB)L_\Delta f, \psi)_\Delta = -(L_\Delta f, \Phi(0, x))_\Delta. \quad (5.30)$$

Since $\Phi(0, x) = B^*\delta(x)$, ($\delta(x)$ is the Dirac function) then according to (5.24) and (5.30) relation (5.27) is true. Let us suppose that in $L(\Delta)$ there is another function $\psi_1(x, s)$ satisfying (5.27). Then the equality

$$((sI - L_\Delta)f, \phi(x, s))_\Delta = 0, \quad \phi = \psi - \psi_1 \quad (5.31)$$

is valid. We write relation (5.31) in the form

$$(L_\Delta f, (I + sB^*)\phi)_\Delta = 0. \quad (5.32)$$

The range of L_Δ is dense in $L^\infty(\Delta)$. Hence in view of (5.32) we have $\phi = 0$. This proves the theorem.

6 The kernel of the quasi-potential

In this section we shall investigate the properties of the kernel $\Phi(x, y)$ of the quasi-potential B .

Proposition 6.1. *Let conditions (5.2), (5.3) be fulfilled and let the quasi-potential B be regular. Then the corresponding kernel $\Phi(x, y)$ is non-negative i.e.*

$$\Phi(x, y) \geq 0. \quad (6.1)$$

Proof. In view of (5.14) and (5.15) we have $\psi(x, s, u) \geq 0$. Relation (5.24) implies that $\psi(x, s) \geq 0$. Now it follows from (5.28) that

$$\Phi(0, x) = \psi(0, x) \geq 0. \quad (6.2)$$

Let us consider the domains Δ_1 and Δ_2 which are connected by relation $\Delta_2 = \Delta_1 + \delta$. We denote the corresponding truncated generators, quasi-potentials kernels by L_k, B_k and $\Phi_k(x, y)$, ($k=1,2$). We introduce the unitary operator

$$Uf = f(x - \delta), \quad (6.3)$$

which maps the space $L^2(\Delta_2)$ onto $L^2(\Delta_1)$. At the beginning we suppose that the conditions of Theorem 2.1 are fulfilled. Using formulas (2.1) and (2.2) we deduce that

$$L_2 = U^{-1}L_1U. \quad (6.4)$$

Hence the equality

$$L_2 = U^{-1}L_1U. \quad (6.5)$$

Hence the equality

$$B_2 = U^{-1}B_1U \quad (6.6)$$

is valid. The last equality can be written in the terms of the kernels

$$\Phi_2(x, y) = \Phi_1(x + \delta, y + \delta). \quad (6.7)$$

According to (6.2) and (6.6) we have

$$\Phi_1(x + \delta, y + \delta) \geq 0. \quad (6.8)$$

As δ is an arbitrary real number relation (6.1) follows directly from (6.7). We remark that an arbitrary generator operator L can be approximated by the operators of form (2.1) (see [], Ch.). Hence the proposition is proved.

In view of (4.1), (4.5) and relation $Bf \in C_\Delta$ the following assertion is true.

Proposition 6.2. *Let the quasi-potential B be regular. Then the equalities*

$$\Phi(a_k, y) = \Phi(b_k, y) = 0 \quad 1 \leq k \leq n \quad (6.9)$$

are valid.

7 Sectorial operators

We introduce the following notions.

Definition 7.1. The bounded operator B in the space $L^2(\Delta)$ is called *sectorial* if

$$(Bf, f) \neq 0, \quad f \neq 0 \quad (7.1)$$

and

$$-\frac{\pi}{2}\beta \leq \arg(Bf, f) \leq \frac{\pi}{2}\beta, \quad 0 \leq \beta \leq 1. \quad (7.2)$$

Definition 7.2. The sectorial operator B is called *strong sectorial* if $\beta < 1$. It is easy to see that the following assertion is true.

Proposition 7.1. *Let the operator B be sectorial. Then the operator $(I + sB)^{-1}$ is bounded when $s \geq 0$.*

References

1. Bertoin J., Levy Processes, University Press, Cambridge, 1996.
2. Ito K., On Stochastic Differential Equations, Memoirs Amer. Math. Soc. No.4, 1951.
3. Kac M., On Some Connections Between Probability Theory and Differential and Integral Equations, Proc. Second Berkeley Symp. Math. Stat. and Prob. Berkeley, 189-215, 1951
4. Kac M., Probability and Related Topics in Physical Sciences, Colorado, 1957.
5. Sakhnovich L.A., Integral Equations with Difference Kernels on Finite Intervals, Operator Theory, v.84, Birkhauser, 1996.
6. Sato K., Levy Processes and Infinitely Divisible Distributions, University Press, Cambridge, 1999.
7. Schoutens W., Symens S., The Pricing of Exotic Options by Monte-Carlo Simulation in a Levy Market with Stochastic Volatility, Preprint, 1-27, 2002
8. Titchmarsh E.C., Introduction to the Theory of Fourier Integrals, Oxford, 1937